

2016

Session C, 2016 First Place: Avian aggression levels in response to the Black-capped Chickadee (*Parus atricapillus*) alarm call in edge habitats around Cranberry Lake Biological station

Audrey Sellepack, Jenna Holakovsky, Alexandra Grove, Joseph Retelskyj

Follow this and additional works at: <http://digitalcommons.esf.edu/clbs>



Part of the [Animal Sciences Commons](#), and the [Forest Sciences Commons](#)

Recommended Citation

Audrey Sellepack, Jenna Holakovsky, Alexandra Grove, Joseph Retelskyj, "Session C, 2016 First Place: Avian aggression levels in response to the Black-capped Chickadee (*Parus atricapillus*) alarm call in edge habitats around Cranberry Lake Biological station" (2016). *Cranberry Lake Biological Station*. Paper 14.

<http://digitalcommons.esf.edu/clbs/14>

This Presentation is brought to you for free and open access by the Environmental and Forest Biology at Digital Commons @ ESF. It has been accepted for inclusion in Cranberry Lake Biological Station by an authorized administrator of Digital Commons @ ESF. For more information, please contact digitalcommons@esf.edu.

Avian aggression levels in response to the Black-capped Chickadee alarm call in edge habitats around Cranberry Lake Biological Station.

Joseph Retelskyj, Alexandra Grove, Audrey Sellepack, & Jenna Holakovsky

Photo by Audrey Sellepack (2016)



State University of New York
College of Environmental Science and Forestry

Background


- Mobbing behavior is an anti-predation strategy used by avian species to drive out a predator spotted in an area (Stewart 1956)
- Behavior includes calling, swooping, and directly hitting the predator (Stewart 1956)
- Alarm calls may differ in acoustic structure depending on situation leading to a more effective defense (Templeton et al 2005)

Rationale

- Is the Black-capped Chickadee essential to eliciting an aggressive mobbing response to potential predators?
- The Black-capped Chickadee may have an important role in protecting a habitat by initiating mobbing (Templeton et al 2005)
- Experimental units: Playback calls
- Sampling units: Quadrates


Playbacks

Black-capped Chickadee
Poecile atricapillus



Black-capped Chickadee is well recognized within avian communities because it is widespread.

Blue Jay
Cyanocitta cristata



Blue Jay participates in mobbing, is a noisy species, and is common throughout North America.

Red-eyed Vireo
Vireo olivaceus



Red-eyed Vireo has a complex song and is another common species throughout North America.

(Cornell 2015)

Hypothesis

- H_{a1} Mobbing bird species in edge habitats will exhibit more occurrences of aggressive behaviors, including high frequency calls and physical contact, in response to Black-capped Chickadee alarm calls than playbacks of Chickadee song, Blue-Jay (*Cyanocitta cristata*), and Red-eyed Vireo (*Vireo olivaceus*) alarm calls.
- H_{o1} There will be no significant difference in occurrences of aggressive behaviors between playbacks.

Hypothesis: Continued


- H_{a2} There will not be significant differences between selected edge habitats as measured by average fly-ins to our quadrates.
- H_{o2} There will be a significant difference between the selected edge habitats.
- H_{a3} There will be a significant difference between the Black-capped Chickadee alarm call and its song.
- H_{o3} There will not be a significant difference between the Black-capped Chickadee alarm call and its song.

Methods: Continued

- 5 meter radius (Doran 2005)
- 5 minute playback time (Doran 2005)



Photo by Jenna Holakovsky (2016)



Black-capped Chickadee



Blue Jay




Red-eyed Vireo



Photos by Jenna Holakovsky (2016)

Audio: Stokes (2010)


Methods: Continued



Methods: Continued



Photos by Jenna
Holakovsky (2016)



Photos by Audrey
Sellepack (2016)



Standard Aggression Index (SAI)

Behavior Name & Code	Rank	Description
Vocalization Low-Frequency (VL)	1	Subject bird is vocalizing at low frequency; indicates low agitation; includes songs and chirps
Vocalization High-Frequency (VH)	3	Subject bird is vocalizing at a high frequency; indicates high agitation; includes alarm calls
Flight Swooping (FS)	4	Subject bird is swooping in direct vicinity of decoy owl; indicates acknowledgment of predator; warning signal; high agitation
Flight In Zone (FZ)	2	Subject bird is in flight within zone; indicates agitation; does not include slight movement from perch to perch
Hit (H)	5	Subject bird has made contact with decoy by means of pecking, scratching, projectile vomiting, projectile defecating; use of body to hit, etc.
Perching (P)	2	Subject bird has entered the proximity and is perching in the zone
Not Visible (NV)	0	Subject bird is not visible within zone

Methods: Continued

Tests: Chi-square and ANOVA

Independent variable: Playback calls

Dependent variable: Aggression levels of birds

Photo by Audrey Sellepack (2016)



Results

Beaver Pond	Forsaith Bog	Old Beaver Pond
BCCH	BCCH	BCCH
BLJA	BLJA	BLJA
BTBW	BTBW	BTBW
CEWA	CEWA	CEWA
COYE	COYE	COYE
OVEN	OVEN	OVEN
PIWO	PIWO	PIWO
REVI	REVI	REVI
WIWR	WIWR	WIWR
WTSP	WTSP	WTSP
BTGW	AMGO	AMRE
CORA	AMRE	BTGW
PIWA	HAWO	CSWA
	KILL	KILL
	MAWA	MAWA
	PIWA	PUFI
	WBNU	YRWA
	YRWA	

- 103 individuals
- 22 different species

Table 1. Alpha codes for 22 different bird species viewed over 3 different edge habitats near Cranberry Lake Biological Station, Clifton, NY. The codes in bold indicate the shared species over the three edge habitats.

Results: ANOVA




Figure 1. Average number of fly-ins in each edge habitat (+/- 1SD).
Cranberry Lake Biological Station, Clifton, New York.

Results: Chi-square




Figure 2. The distribution of observed aggression through four quadrant ranges of playbacks across a total of 12 quadrates (n=48). Cranberry Lake Biological Station, Clifton, New York.



Discussion:

We reject the null for H_{a1}

- Black-capped Chickadees had highest aggression ranks
- Discrepancies in Data

We reject the null for H_{a2}

- No difference between edge habitats

We reject the null for H_{a3}

- Significant difference between call and song

Discussion:

- Bird communities coordinate anti-predatory behavior through interspecific communication (Hurd 1996)
- Local birds respond to alarm calls of *Paridae* species rather than non-*Paridae* species (Langham et al 2009)
- Birds will exhibit a wide range of mobbing behaviors around a potential predator (Curio 1978)

Photo by Audrey Sellepack (2016)



Discussion:

Methodology Drawbacks

- Larger quadrate circumference
- Additional edge habitat sites
- Further isolate transects

Photo by Audrey Sellepack (2016)



Discussion:

Potential Follow-up Studies

- Repeat study in different habitats
- Play *Paridae* alarm calls, including different species
- Black-capped Chickadees and mobbing bird mortality rates

Photo by Jenna Holakovsky (2016)



Conclusion:

- Mobbing bird species tended to behave more aggressively when a Black-capped Chickadee alarm call was played rather than a Blue Jay alarm call or Red-eyed Vireo alarm call
- Difference between Black-capped Chickadee alarm call and the Black-capped Chickadee song
- No differences between selected edge habitats

Acknowledgements

- The Cranberry Lake Biological Station
- SUNY ESF and Dr. Fierke
- Ornithologist, Alan Belford
- Thomas Evans
- Giuseppe Tumminello
- Roxie and kitchen staff
- Indiana State University White-throated Sparrow Research Team
- A special thank you to the bear for checking in on us and keeping us awake.

Photo by Tyler Hodges (2016)



References

- Altmann, Stuart. "Avian Mobbing Behavior and Predator Recognition." *The Condor* 58.4 (1956): 241-53. *JSTOR*. Web. 14 July 2016.
- Curio, Eberhard. "The adaptive significance of avian mobbing I: teleonomic hypotheses and predictions." *Zeitschrift fur Tierpsychologie* 48.2 (1978): 175-83. *PsycINFO*. Web. 14 July 2016.
- Doran, Patrick J., Paul Z. Gulezian, and Matthew G. Betts. "A Test of the Mobbing Playback Method for Estimating Bird Reproductive Success." *Journal of Field Ornithology* 76.3 (2005): 227-33. *JSTOR*. Web. 7 July 2016.
- Hurd, Christine R. "Interspecific Attraction to the Mobbing Calls of Black-Capped Chickadees (*Parus atricapillus*)." *Behavioral Ecology and Sociobiology* 38.4 (1996): 287-92. *JSTOR*. Web. 7 July 2016.
- Langham, Gary M., Thomas A. Contreras, and Katheryn E. Sieving. "Why pishing works: Titmouse (*Paridae*) scolds elicit a generalized response in bird communities." *Ecoscience* 13.4 (2006): 485-96. *BioOne*. Web. 7 July 2016.
- McPherson, Roger, and Richard Brown. "Mobbing Responses of some Passerines to the Calls and Location of the Screech Owl." *Raptor Research* 15.1 (1981): 23-30. *SORA*. Web. 14 July 2016.
- Rice, Jake C. "Behavioral Implications of Aberrant Song of a Red-Eyed Vireo." *Wilson Bulletin* 93.3 (1981): 383-90. *jstor*. Web. 14 July 2016.
- Templeton, Christopher, Erick Greene, and Kate Davis. "Allometry of Alarm Calls: Black-Capped Chickadees Encode Information About Predator Size." *Science* 308.5730 (2005): n. pag. *Science*. Web. 14 July 2016.

Questions?

Photo by Jenna Holakovsky (2016)

